arXiv:hep-ex/0605055v1 15 May 2006

KAON PHYSICS AT KLOE

THE KLOE COLLABORATION[a]
presented by MATTEO PALUTAN
INFN, Laboratori Nazionali di Frascati, via E. Fermi 40, 00044 - Frascati, Italy

The most precise determination of $V_{\rm us}$ comes from semileptonic kaon decays. We have measured with the KLOE detector at DAΦNE, the Frascati ϕ factory, all the experimental inputs to $V_{\rm us}$ for both neutral and charged kaons. Using our results we extract the value of $V_{\rm us}$ with $\sim 0.9\%$ fractional error, which is entirely dominated by the theoretical error on the knowledge of the normalization of the decay form factors, $f_+(0)$. A new determination of the ratio $V_{\rm us}/V_{\rm ud}$ is also presented, based on our precise measurement of the absolute branching ratio for the decay $K \to \mu\nu$, combined with recent lattice results for the ratio f_K/f_π. New results on CPT symmetry have also been achieved, which are based on the first measurement of the charge asymmetry for $K_S \to \pi e \nu$ decay.

1 Introduction

The CKM matrix element $V_{\rm us}$ can be extracted from the measurement of the semileptonic decay widths ($\Gamma = {\rm BR}/\tau$) and the most precise test of unitarity of the CKM matrix is performed from the first-row constraint: $1-\Delta = |V_{\rm ud}|^2 + |V_{\rm us}|^2 + |V_{\rm ub}|^2$. Using $V_{\rm ud}$ from nuclear beta decays, a test of the expectation $\Delta = 0$ with a precision of one part per mil can be performed, $V_{\rm ub}$ contributing only at the level of 10^{-5}. With the KLOE detector we can measure all experimental inputs to $V_{\rm us}$: branching ratios, lifetimes, and decay form factors.

---

[a] F. Ambrosino, A. Antonelli, M. Antonelli, C. Bacci, P. Beltrame, G. Bencivenni, S. Bertolucci, C. Bini, C. Bloise, V. Bocci, F. Bossi, D. Bowring, P. Branchini, R. Caloi, P. Campana, G. Capon, T. Capussela, F. Ceradini, S. Chi, G. Chiefari, P. Ciambrone, S. Conetti, E. De Lucia, A. De Santis, P. De Simone, G. De Zorzi, S. Dell'Agnello, A. Denig, A. Di Domenico, C. Di Donato, S. Di Falco, B. Di Micco, A. Doria, M. Dreucci, G. Felici, A. Ferrari, M. L. Ferrer, G. Finocchiaro, S. Fiore, C. Forti, P. Franzini, C. Gatti, P. Gauzzi, S. Giovannella, E. Gorini, E. Graziani, M. Incagli, W. Kluge, V. Kulikov, F. Lacava, G. Lanfranchi, J. Lee-Franzini, D. Leone, M. Martini, P. Massarotti, W. Mei, S. Meola, S. Miscetti, M. Moulson, S. Müller, F. Murtas, M. Napolitano, F. Nguyen, M. Palutan, E. Pasqualucci, A. Passeri, V. Patera, F. Perfetto, L. Pontecorvo, M. Primavera, P. Santangelo, E. Santovetti, G. Saracino, B. Sciascia, A. Sciubba, F. Scuri, I. Sfiligoi, T. Spadaro, M. Testa, L. Tortora, P. Valente, B. Valeriani, G. Venanzoni, S. Veneziano, A. Ventura, R.Versaci, G. Xu

In the DAΦNE e^+e^- collider, beams collide at a center-of-mass energy $W \sim M(\phi)$. Since 2001, KLOE has collected an integrated luminosity of $\sim 2500\,\mathrm{pb}^{-1}$. Results presented below are based on 2001-2002 data, corresponding to an integrated luminosity of $\sim 450\,\mathrm{pb}^{-1}$. The KLOE detector consists of a large cylindrical drift chamber surrounded by a lead/scintillating-fiber electromagnetic calorimeter. A superconducting coil around the calorimeter provides a 0.52 T field. The drift chamber, [1] is 4 m in diameter and 3.3 m long. The momentum resolution is $\sigma(p_T)/p_T \sim 0.4\%$. Two track vertices are reconstructed with a spatial resolution of ~ 3 mm. The calorimeter, [2] composed of a barrel and two endcaps, covers 98% of the solid angle. Energy and time resolution are $\sigma(E)/E = 5.7\%/\sqrt{E(\,\mathrm{GeV})}$ and $\sigma(t) = 57\ \mathrm{ps}/\sqrt{E(\,\mathrm{GeV})} \oplus 100\ \mathrm{ps}$. The KLOE trigger, [3] uses calorimeter and drift chamber information. For the present analysis only the calorimeter signals are used. Two energy deposits above threshold, $E > 50$ MeV for the barrel and $E > 150$ MeV for the endcaps, are required.

The ϕ meson decays mainly into kaons: 49% to $K^+K^-$and 34% to K_SK_L pairs. We can thus tag K_L, K_S, K^+, and K^- decays by detecting respectively K_S, K_L, K^-, and K^+ decays on the opposite hemisphere. The tagging technique is a unique opportunity of a ϕ factory, and provides pure and almost monochromatic "beams" of kaons, thus allowing to measure the absolute branching ratios. The K_S beam is tagged using events with a K_L interacting in the calorimeter (K_L-crash). K_L-mesons are tagged detecting $K_S \to \pi^+\pi^-$ decays. Charged kaons are tagged using two-body decays, $K^\pm \to \mu^\pm\nu$ and $K^\pm \to \pi^\pm\pi^0$. For all of the cases it is possible to precisely measure the tagged kaon momentum from the knowledge of the ϕ and the tagging kaon momenta.

2 K_L decays

2.1 Major K_L decays and lifetime

As already stated, a pure sample of K_L decays is selected by the identification of $K_S \to \pi^+\pi^-$ decays. Events include K_L which decay to any possible final state in the detector volume, interact in the calorimeter or escape the detector. Starting from this sample, the K_L branching ratios are evaluated by counting the number of decays to each channel in the fiducial volume (FV) and correcting for the geometrical acceptance, the reconstruction efficiency, and the background contamination.

K_L charged decays are identified by selecting a decay vertex in the FV matching the expected K_L flight direction, as defined by the tag. In order to discriminate among the different K_L charged modes the variable $\Delta_{\mu\pi} = |p_{miss} - E_{miss}|$ is used, where p_{miss} and E_{miss} are the missing momentum and missing energy at the K_L vertex, evaluated by assigning one track the pion mass and the other one the muon mass (left panel of Fig. 1). Signal counting is thus achieved by fitting the $\Delta_{\mu\pi}$ spectrum with a linear combination of four Monte Carlo shapes ($K_L \to \pi e\nu, \pi\mu\nu, \pi^+\pi^-\pi^0, \pi^+\pi^-$). The shape of $\Delta_{\mu\pi}$ spectrum is sensitive to the radiative corrections to K_L decay processes. The KLOE Monte Carlo includes very accurate generators for all of the radiative channels, which has been studied and implemented mostly in 2004. [4].

To count $K_L \to \pi^0\pi^0\pi^0$ events decaying in the FV, we exploit the time-of-flight capability of the calorimeter to reconstruct the neutral vertex position. Such a vertex is assumed to be along the K_L line of flight. The arrival time of each photon identified in the calorimeter is thus used to give an independent determination of the path length of the K_L, L_K. The final value of L_K is obtained from a wheighted average of the different measurements coming from all of the photons. The analysis of the $K_L \to \pi^0\pi^0\pi^0$ sample is used not only to count events for BR measurement, but also to extract the K_L lifetime (τ_{K_L}) from a fit to the proper time distribution of neutral decay vertices. The KLOE environment is indeed very well suited for a direct measurement of τ_{K_L}, because of the tagging technique which provides the value of the K_L momentum and allows



Figure 1: Left panel: distribution of $\Delta_{\mu\pi}$ for a data subsample, with fit to MC distributions for different decay channels. Right panel: KLOE results on semileptonic form factor slopes compared with other recent measurements; the black dot represents the values of λ'_+ and λ''_+ obtained from the Taylor expansion of the pole parametrization.

to select a pure sample of K_L decays. The result is $\tau_{K_L} = 50.92 \pm 0.17 \pm 0.25$ ns. [5]

A total of about 13×10^6 tagged events have been used for the measurement of the K_L BR's. Since the geometrical efficiency of the fiducial volume depends on the value of the K_L lifetime, the sum of all of the decay modes is fixed to unity (BR's below 1% from PDG [11]). This removes the uncertainty due to τ_{K_L}, while giving at the same time a precise determination of K_L lifetime itself. Results for the main BR's are: [6]

$$\begin{array}{rcl} \mathrm{BR}(K_L \to \pi e \nu) & = & 0.4007 \pm 0.0005_{\mathrm{stat}} \pm 0.0004_{\mathrm{syst-stat}} \pm 0.0014_{\mathrm{syst}}\,, \\ \mathrm{BR}(K_L \to \pi \mu \nu) & = & 0.2698 \pm 0.0005_{\mathrm{stat}} \pm 0.0004_{\mathrm{syst-stat}} \pm 0.0014_{\mathrm{syst}}\,, \\ \mathrm{BR}(K_L \to \pi^0\pi^0\pi^0) & = & 0.1997 \pm 0.0003_{\mathrm{stat}} \pm 0.0004_{\mathrm{syst-stat}} \pm 0.0019_{\mathrm{syst}}\,, \\ \mathrm{BR}(K_L \to \pi^+\pi^-\pi^0) & = & 0.1263 \pm 0.0004_{\mathrm{stat}} \pm 0.0003_{\mathrm{syst-stat}} \pm 0.0011_{\mathrm{syst}}\,. \end{array} \tag{1}$$

The corresponding lifetime is $\tau_{K_L} = 50.72 \pm 0.11_{\mathrm{stat}} \pm 0.13_{\mathrm{syst-stat}} \pm 0.33_{\mathrm{syst}}$ ns, in agreement with the direct measurement that has been shown before. The two lifetime measurements are uncorrelated; their average is $\tau_{K_L} = 50.84 \pm 0.23\,\mathrm{ns}$, and has a fractional error of 0.45%. This represents a factor of two improvement with respect to the previous measurements. [7]

2.2 Semileptonic form factor slopes

In semileptonic kaon decays, $K_{L,S} \to \pi^\pm \ell^\mp \nu$, only the vector part of the weak current has a non-vanishing matrix element between a kaon and a pion. The vector current is "almost" conserved. For a vector interaction, there are no $SU(3)$-breaking corrections to first order in the s-d mass difference, making calculations of hadronic matrix elements more reliable. In the electron mode $K_{L,S} \to \pi^\pm e^\mp \nu$, only the vector form factor $f_+(t)$ is involved, since extra terms in the matrix element depend on the electron mass. This form factor is usually parametrized as

$$f_+(t) = f_+(0) \left[1 + \lambda'_+ \frac{t}{m^2_{\pi^+}} + \frac{\lambda''_+}{2} \frac{t^2}{m^4_{\pi^+}} + \ldots \right], \tag{2}$$

where $f_+(0)$ is evaluated from theory and t is the lepton-pair invariant mass squared. We used about 2 million of $K_L \to \pi e \nu$ decays from 2001-2002 data sample to measure the semileptonic form factor slopes. These have been extracted by fitting the spectrum of $t/m^2_{\pi^+}$ for the selected events. The fit procedure takes into account the efficiency of the selection cuts, the resolution effects and the background contamination as a function of t. The results obtained for a quadratic fit are:[10]

$$\begin{array}{rcl} \lambda'_{+} & = & (25.5 \pm 1.5_{\rm stat} \pm 1.0_{\rm syst}) \times 10^{-3} \\ \lambda''_{+} & = & (1.4 \pm 0.7_{\rm stat} \pm 0.4_{\rm syst}) \times 10^{-3}, \end{array} \tag{3}$$

which gives a $\chi^2/ndf = 325/362$ $(P(\chi^2) = 92\%)$.

We have also fit the data using a pole parametrization: $f_+(t)/f_+(0) = M_V^2/(M_V^2 - t)$. This assumes the form factor is dominated by the vector $K - \pi$ resonances, the closest being the $K^*(892)$. We obtain $M_V = (870 \pm 6_{\rm stat} \pm 7_{\rm syst})$ MeV $(\chi^2/ndf = 326/363)$, that confirms the dominance of K^* meson, although contributions from other $J^P = 1^-$ resonant and non-resonant $K\pi$ scattering amplitudes are not negligible. Note that the quadratic parametrization of the form factor arises naturally as a Taylor expansion of the pole model, with the additional contraint $\lambda''_+ = 2\lambda_+^2$. This is nicely fullfilled by our result, which is competitive with other recent measurements[8, 9] (right panel of Fig. 1).

3 $K_S \to \pi e \nu$ decay

At the ϕ factory very large samples of tagged, monochromatic K_S mesons are available. From the analysis of 2001-2002 data sample, we were able to obtain a very pure sample of $\sim 13\,000$ K_S semileptonic decay events and thus to accurately measure for the first time the partial decay rates for transitions to each charged final state, BR$(K_S \to e^+\pi^-\nu)$ and BR$(K_S \to e^-\pi^+\bar{\nu})$, as well as the charge asymmetry A_S:

$$A_S = \frac{\Gamma\,(\mathrm{K_S} \to \pi^-\mathrm{e}^+\nu) - \Gamma\,(\mathrm{K_S} \to \pi^+\mathrm{e}^-\bar{\nu})}{\Gamma\,(\mathrm{K_S} \to \pi^-\mathrm{e}^+\nu) + \Gamma\,(\mathrm{K_S} \to \pi^+\mathrm{e}^-\bar{\nu})}. \tag{4}$$

The comparison of A_S with the corresponding asymmetry A_L for K_L decays allows precision tests of the CP and CPT symmetries. If CPT symmetry is assumed, both K_S and K_L charge asymmetries are expected to be equal to $2\,\mathrm{Re}\,\epsilon \simeq 3 \times 10^{-3}$. The difference between the charge asymmetries,

$$A_S - A_L = 4\;(\mathrm{Re}\,\delta + \mathrm{Re}\,x_-)\,, \tag{5}$$

signals CPT violation either in the mass matrix (δ term), or in the decay amplitudes with $\Delta S \neq \Delta Q$ (x_- term). The sum of the asymmetries,

$$A_S + A_L = 4\,(\mathrm{Re}\,\epsilon - \mathrm{Re}\,y)\,, \tag{6}$$

is related to CP violation in the mass matrix (ϵ term) and to CPT violation in the decay amplitude (y term). The knowledge of both the K_L and the K_S semileptonic decay branching ratios and lifetimes allows the validity of the $\Delta S = \Delta Q$ rule to be tested through the quantity

$$\mathrm{Re}\,x_+ = \frac{1}{2}\frac{\Gamma(K_S \to \pi e\nu) - \Gamma(K_L \to \pi e\nu)}{\Gamma(K_S \to \pi e\nu) + \Gamma(K_L \to \pi e\nu)}. \tag{7}$$

In the SM, $\mathrm{Re}\,x_+$ is of the order of $G_F m_\pi^2 \sim 10^{-7}$, being due to second order weak transitions. Finally, from the semileptonic decays of K_S a competitive measurement of $V_{\rm us}$ can be extracted, which also profits of a very precise knowledge of the K_S lifetime.[11]

The first measurement of BR$(K_S \to \pi e \nu)$ was obtained from KLOE using data collected in 2000, with a fractional accuracy of 5.4%.[12] The present BR measurement improves on the total error by a factor of four, to 1.3%. The K_S charge asymmetry has never been measured before. Counting of events is performed by fitting the $E_{miss} - p_{miss}$ spectrum with a combination of MC shapes for signal and background (Fig. 2). For each charge state we obtain the ratio BR$(K_S \to \pi^\mp e^\pm \nu(\bar{\nu}))$/BR$(K_S \to \pi^+\pi^-)$ by normalizing the number of signal events to the



Figure 2: $E_{miss} - p_{miss}$ spectrum for events selected as $\pi^- e^+ \nu$ (left panel) and $\pi^+ e^- \overline{\nu}$ (right panel); filled dots represent data from the entire data set; solid line is the result of a fit varying the normalization of MC distributions for signal (light gray) and background (dark gray), which are also shown.

number of $K_S \to \pi^+\pi^-$ events and correcting for the overall selection efficiencies. In order to evaluate the BR's for the semileptonic modes, we combine the measured ratios with the ratio R_S^π of the dominant K_S decay modes which is measured at KLOE. We obtain the following results:[16]

$$\begin{array}{rcl} \mathrm{BR}(K_S \to \pi^- e^+ \nu) & = & (3.528 \pm 0.062) \times 10^{-4} \\ \mathrm{BR}(K_S \to \pi^+ e^- \bar{\nu}) & = & (3.517 \pm 0.058) \times 10^{-4} \\ \mathrm{BR}(K_S \to \pi e \nu) & = & (7.046 \pm 0.091) \times 10^{-4}. \end{array} \tag{8}$$

The charge asymmetry of Eq. (4) is evaluated to be:[16] $A_S = (1.5 \pm 9.6_{\rm stat} \pm 2.9_{\rm syst}) \times 10^{-3}$. From the total BR we test the validity of the $\Delta S = \Delta Q$ rule in CPT-conserving transitions (Eq. 7). In the evaluation of $\Gamma's$ we used $\tau_{K_S} = 0.08958(6)$ ns from the PDG[11] and KLOE results for τ_{K_L} and $\mathrm{BR}(K_L \to \pi e \nu)$. We obtain $\mathrm{Re}\, x_+ = (-0.5 \pm 3.6) \times 10^{-3}$.[16] The error on this measurement represents an improvement by almost a factor of two with respect to the most precise previous result.[13]

From the sum and difference of the K_L and K_S charge asymmetries one can test for possible violations of the CPT symmetry, either in the decay amplitudes or in the mass matrix (Eqs. 5 and 6). Using $A_L = (3.34 \pm 0.07) \times 10^{-3}$,[11] and $\mathrm{Re}\,\delta = (3.0 \pm 3.3_{\rm stat} \pm 0.3_{\rm syst}) \times 10^{-4}$,[14] we obtain $\mathrm{Re}\, x_- = (-0.8 \pm 2.5) \times 10^{-3}$.[16] This result improves by a factor of ten respect to the previous best result.[14] Using $\mathrm{Re}\,\epsilon = (1.62 \pm 0.04) \times 10^{-3}$,[11] we obtain $\mathrm{Re}\, y = (0.4 \pm 2.5) \times 10^{-3}$,[16] which has precision comparable to that (3×10^{-3}) obtained from the unitarity relation by CPLEAR.[15]

Finally, we measured the form factor slope by fitting the ratio of data and MC distributions in $t/m^2_{\pi^+}$. The fit has been performed by using only a linear parametrization of the form factor, since the available statistics do not allow to be sensitive to a quadratic one. The linear slope of the semileptonic K_S form factor is measured to be $\lambda_+ = (33.9 \pm 4.1) \times 10^{-3}$.[16] This result is in agreement with the corresponding value for the linear slope of the semileptonic K_L form-factor.

4 Charged kaon decays

4.1 Semileptonic $K^\pm$ decays

The measurement of the branching ratios for the $K^\pm$ semileptonic decays is performed using four data samples defined by different decay modes and charge states of the tagging kaon: $K^+_{\mu 2}$,

$K^+_{\pi 2}$, $K^-_{\mu 2}$, and $K^-_{\pi 2}$. This redundancy allows to keep under control the systematic effects due to the tag selection. Kaons are identified as tracks with momentum $70 < \mathrm{p} < 130\,\mathrm{MeV}$, originating from the collision point. The kaon decay vertex must be within a fiducial volume (FV) defined as a cylinder of radius $40 < \mathrm{r} < 150\,\mathrm{cm}$, centered at the collision point, coaxial with the beams. The decay track, extrapolated to the calorimeter, must point to an appropriate energy deposit. $K_{\mu 2}$ ($K_{\pi 2}$) decays are selected by applying a 3σ cuts around the muon (pion) momentum calculated in the kaon rest frame, according to the proper mass hypothesis. For the $K^{\pm}_{\pi 2}$ tag, identification of the π^0 from the vertex is also required. Finally, to reduce the dependence of the tag selection efficiency on the decay mode of the signal kaon, the tagging decay is required to satisfy the calorimeter trigger by itself. To select a semileptonic decay on the signal side, a one-prong kaon decay vertex must be present in the FV. The daughter track has to reach the calorimeter and to overlap an energy deposit; two-body decays are rejected by requiring that its momentum in the kaon frame, computed assuming the pion mass, is less than 195 MeV. The lepton mass (m_{lept}) is obtained from the velocity of the lepton computed from the time of flight. The number of K_{e3} and $K_{\mu 3}$ decays is then obtained by fitting the m^2_{lept} distribution (Fig. 3) to a sum of MC distributions for the signals and the various background sources.



Figure 3: Lepton mass distribution of the K^-_{l3} sample tagged by $K^+_{\mu 2}$ events; K_{e3} events are recognized as a peak around $m^2_{lept} = 0$, while $K_{\mu 3}$ events peak around $m^2_{lept} \sim m^2_{\mu}$.

The BR is evaluated separately for each tag sample, dividing the number of signal counts by the number of tag events and correcting for acceptances. The latter are obtained from MC simulation, and correcting for data-MC differences in tracking efficiency and calorimeter clustering. About 190 000 $K^{\pm}_{e3}$ and 100 000 $K^{\pm}_{\mu 3}$ decays are selected. Preliminary results for the BR's are:

$$\begin{aligned} \mathrm{BR}(K^{\pm}_{e3}) &= (5.047 \pm 0.046_{\mathrm{stat}} \pm 0.080_{\mathrm{syst}})\% \\ \mathrm{BR}(K^{\pm}_{\mu 3}) &= (3.310 \pm 0.040_{\mathrm{stat}} \pm 0.070_{\mathrm{syst}})\%. \end{aligned} \tag{9}$$

These values have been averaged over the four different tag samples for each channel. The quoted systematic error is still preliminary.

4.2 $K^{\pm}$ lifetime

The $K^{\pm}$ lifetime ($\tau_{K^{\pm}}$) is an experimental input to the determination of V_{us}. The present fractional uncertainty is about 0.2%, corresponding to an uncertainty of 0.1% for V_{us}. However, there are large discrepancies between results of different experiments.[11] The value of $\tau_{K^{\pm}}$ also affects the geometrical acceptance in BR measurements. KLOE is measuring $\tau_{K^\pm}$ with high-statistics using two different methods: one based on the measurement of the decay length and the other on the decay time of the kaons. Comparison between the two methods will allow to decrease the systematic error. The preliminary result from the decay length measurement is $\tau_{K^\pm} = 12.336 \pm 0.044_{\rm stat} \pm 0.065_{\rm syst}$ ns.

5 Determination of $V_{\rm us}$

$V_{\rm us}$ is proportional to the square root of the partial width of semileptonic kaon decays , and can be parametrized, for neutral kaon decays, as:

$$|V_{\rm us} f_+(0)| = \left[\frac{128\pi^3\Gamma}{G_\mu^2 M_K^5 S_{\rm ew} I(\lambda'_+, \lambda''_+, \lambda_0, 0)}\right]^{1/2} \frac{1}{1+\delta_{\rm em}}, \tag{10}$$

where $f_+(0)$ is the vector form factor at zero momentum transfer and $I(\lambda'_+, \lambda''_+, \lambda_0, 0)$ is the result of the phase space integration after factorizing out $f_+(0)$. In the above expression, long-distance radiative corrections for both the form factor $f_+(0)$ and the phase space integral have been factorized out and are included in the parameter $\delta_{\rm em}$, which amounts to $5 - 8 \times 10^{-3}$ for K_{e3} and $K_{\mu3}$ respectively.[17, 18] The short-distance electroweak corrections are included in the parameter $S_{\rm ew} = 1.0232$;[19] λ'_+ and λ''_+ are the quadratic slopes of the vector form factor, and λ_0 is the slope of the scalar form factor.

The BR's of the semileptonic K_L decays (and τ_{K_L}), together with the measurement of BR($K_S \to \pi e \nu$) and the preliminary results on the semileptonic $K^\pm$ decays, give five independent determinations of the observable $|V_{\rm us} f_+(0)|$. The average of KLOE results is $|V_{\rm us} f_+(0)| = 0.2166 \pm 0.0005$, with $\chi^2/ndf = 1.9/4$. The fractional uncertainty is about 0.25%. A precise estimate of $f_+(0) = 0.961 \pm 0.008$ was first given in 1984,[20] and it has been recently confirmed by lattice calculations.[21] Using the previous value for $f_+(0)$ and the KLOE result for $|V_{\rm us} f_+(0)|$ we find $V_{\rm us} = 0.2254 \pm 0.0020$. This has to be compared with the value obtained from unitarity contraint: $V_{\rm us} = \sqrt{1 - V_{\rm ud}^2} = 0.2275 \pm 0.0012$.[22]

A complementary approach to the previous determination of $V_{\rm us}$ is the extraction of $|V_{\rm us}|/|V_{\rm ud}|$ from the measurement of the ratio of kaon and pion leptonic decay rates, which is parametrized by the equation

$$\frac{\Gamma(K \to \mu\nu)}{\Gamma(\pi^{)} \to \mu^{)}\nu} = \frac{|V_{\rm us}|^2}{|V_{\rm ud}|^2} \frac{m_K \left(1 - \frac{m_\mu^2}{m_K^2}\right)^2}{m_\pi \left(1 - \frac{m_\mu^2}{m_\pi^2}\right)^2} \frac{f_K^2}{f_\pi^2} \frac{1 + \frac{\alpha}{\pi} C_K}{1 + \frac{\alpha}{\pi} C_\pi}, \tag{11}$$

where f_K and f_π are the kaon and the pion decay contants, respectively; C_K and C_π parametrize the radiative electroweak corrections. KLOE has recently measured the absolute $K^+_{\mu2}$ branching ratio, with inclusion of photon radiation in the final state: BR($K^+ \to \mu^+\nu$)=$0.6366 \pm 0.0009_{\rm stat}$ $\pm 0.0015_{\rm syst}$.[23] Using a recent lattice calculation of the ratio f_K/f_π of pseudoscalar meson decay constants,[24] we find $V_{\rm us}/V_{\rm ud} = 0.2294 \pm 0.0026$. The fractional error is $\sim 1\%$, and is dominated by the uncertainty in the f_K/f_π ratio. The previous results on $V_{\rm us}$ and $V_{\rm us}/V_{\rm ud}$, together with the present knowledge of $V_{\rm ud}$ are shown in Fig. 4; a black line represents the unitarity constraint. A combined fit of the experimental results gives $V_{\rm us} = 0.2246 \pm 0.0016$. In the same $V_{\rm us}$-$V_{\rm ud}$ plane, the experimental data match the unitarity constraint with P(χ^2)$\sim$0.23.

References

1. M. Adinolfi *et al.*, KLOE Collaboration, *Nucl. Instrum. Methods* A **488**, 51 (2002)
2. M. Adinolfi *et al.*, KLOE Collaboration, *Nucl. Instrum. Methods* A **482**, 364 (2002)



Figure 4: Pictorial view of $V_{\rm us}$, $V_{\rm ud}$and $V_{\rm us}/V_{\rm ud}$ measurements in the $V_{\rm us}$ vs $V_{\rm ud}$ plane. The small ellipse represents the fit result for $V_{\rm us}$ and $V_{\rm ud}$; the unitariry contraint is represented by the black line.

3. M. Adinolfi *et al.*, KLOE Collaboration, *Nucl. Instrum. Methods* A **492**, 134 (2002)
4. C.Gatti, *Eur. Phys. J.* C **45**, 417 (2005)
5. KLOE Collaboration, F. Ambrosino *et al.*, *Phys. Lett.* B **626**, 15 (2006)
6. KLOE Collaboration, F. Ambrosino *et al.*, *Phys. Lett.* B **632**, 43 (2006)
7. K.G. Vosburgh *et al.*, *Phys. Rev.* D **6**, 1834 (1972)
8. KTeV Collaboration, T. Alexopoulos *et al.*, *Phys. Rev. Lett.* **93**, 181802 (2004)
9. ISTRA+ Collaboration, O. P. Yushchenko *et al.*, *Phys. Lett.* B **589**, 111 (2004)
10. KLOE Collaboration, F. Ambrosino *et al.*, *Phys. Lett.* B **636**, 166 (2006)
11. Particle Data Group, S. Eidelman *et al.*, *Phys. Lett.* B **592**, 1 (2004)
12. KLOE Collaboration, A. Aloisio *et al.*, *Phys. Lett.* B **535**, 37 (2002)
13. CPLEAR Collaboration, Phys. Lett. B444 (1998) 38.
14. CPLEAR Collaboration, Phys. Lett. B444 (1998) 52.
15. CPLEAR Collaboration, Phys. Lett. B456 (1999) 297.
16. KLOE Collaboration, F. Ambrosino *et al.*, *Phys. Lett.* B **636**, 173 (2006)
17. V. Cirigliano, H. Neufeld, H. Pichl, *Eur. Phys. J.* C **35**, 53 (2004)
18. T. C. Andre, Radiative corrections to K0(l3) decays, **hep-ph/0406006**.
19. A. Sirlin, *Nucl. Phys.* B **196**, 83 (1982)
20. H. Leutwyler, M. Roos, ZPC **25**, 91 (1984)
21. D. Becirevic *et al.*, *Nucl. Phys.* B **705**, 339 (2005)
22. W. J. Marciano, A. Sirlin, *Phys. Rev. Lett.* **96**, 032002 (2006)
23. KLOE Collaboration, F. Ambrosino *et al.*, *Phys. Lett.* B **632**, 76 (2006)
24. MILC Collaboration, C. Bernard *et al.*, “Update on π and K physics”, Presented at 23rd International Symposium on Lattice Field Field: Lattice 2005, Trinity College, Dublin, Ireland, 25-30 Jul 2005. Published in PoS LAT2005:025,2005.